



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**XVII.** *A Letter from Monsieur Wargentin, Secretary to the Royal Academy of Sciences in Sweden, to Mr. John Ellicott, F.R.S. relating to the late Transit of Venus.*

Stockholm, December 24, 1762.

Read March 17, 1763. **T**HE observations upon the last transit of Venus over the Sun made at the Cape of Good Hope are excellent, and seem to decide, that the horizontal parallax of the Sun is  $8''$ , 1 or  $8''$ , 3 at most. I had before found it to be that quantity, from the observations made in Europe compared together; but the observations made at the Cape confirm it with the greatest evidence.

It is of importance to be assured of the longitude of the places where the observations were made. I have endeavoured to determine them the best I was able by observations of the eclipses of Jupiter's satellites, made at the same places. That you may the better judge, I thought proper to send you all the observations of these satellites made at different places the last year. I desire you would communicate them to Mr. Maſon, and all those who interest themselves in the research of the parallax. It is pity that messieurs L'Abbè Chappe, and Rumoski, did not succeed in observing several eclipses of the satellites, at Tobieske and Selenginsk, the better to confirm the Longitudes of those places. However it appears to me, that the difference between the meridians of

Greenwich and Tobieské is scarcely more than  $4^h 32' 55''$ . That between the meridians of Greenwich and Selenginsk, to judge from the three immersions observed there, should be but  $7^h 6' 0$ , but from other considerations, I think it must be 10 or 15' more. If the longitude of these places should be more exactly determined, I am persuaded that we should obtain the parallax of the Sun to nearly the tenth of a second, so exact the observations made at Selenginsk and Tobieské and the Cape appear to me.

I see you did not communicate to the Royal Society all the observations made by Mr. Planman at Cajaneberg, they however deserve to be preserved. He observed

	h	'	''
The beginning of the entrance - - at	3	59	56M
Total immersion of Venus or interior contact - - - - - }	4	18	5
Second interior contact or beginning of the exit - - - - - }	10	7	59
Total emerfion - - - - -	10	26	22

Mr. Planman made use of a telescope of 20 or 21 feet; the latitude of Cajanebourg is  $64^{\circ} 13' 30''$ ; the difference of meridians between Greenwich and Cajaneburg is sufficiently determined by observations upon the eclipse of the Moon the 18th May 1761, made at Stockholm and Cajaneburg.

I have the honour to be with the greatest esteem,

S I R,

Your very humble and obedient servant,

P. Wargentín.

Eclipses primi satellitis Jovis, observatæ Anno 1761 :  
addito errore calculi pro quavis observatione, ut ap-  
pareat, quæ sint præstantiores. In calculo autem  
posui differentiam meridianorum inter

		h	'	"
Observatorium	Grenovic. et Paris.	—	0	9 10
	Grenov. et Stockholm	—	1	12 0
	Grenov. et Upsaliense	—	1	10 20
	Grenov. et Massil.	—	0	21 19
	Grenov. et Cap. B. Spei	—	1	13 35
	Grenov. et Insul. Rodrig.	—	4	12 52
	Grenov. et Selenginsk	—	7	6 15
	Grenov. et Cajaneburgh	—	1	50 40

		h	'	"			
Jan. 7.	Em.	5	4	35	Stockholmia	+	0 27
June 13.	Im.	14	37	47	Massilia - - -	—	0 21*
	20. —	17	22	26	Cap. Bonæ Spei	+	0 31
July 22.	—	12	45	15	Paris - - - -	+	0 7
		12	35	39	Grenov. - - -	+	0 33
		16	48	55	Inf. Rodriga -	+	0 9
29.	—	14	38	35	Paris - - - -	+	0 33
		14	50	28	Massilia - - -	+	0 49
		15	42	44	Cap. B. Spei -	+	0 49
31.	—	13	10	29	Inf. Rodrig. -	+	0 56*
Aug. 7.	—	10	51	52	Grenovic. - - -	+	1 8*
		12	4	51	Stockholm. bona	+	0 9
		11	13	49	Massilia - - -	+	0 30
		12	5	46	Cap. B. Spei -	+	0 49
14.	—	13	59	39	Stockholm. certa.	+	0 10
		14	0	50	Cap. B. Sp. -	+	0 34

Aug.

Aug. 21. Im.	<sup>h</sup>	<sup>'</sup>	<sup>"</sup>								
	14	51	56	Paris	-	-	-	-	+	0	34
	15	56	2	Cap. B. Spei	-	-	-	-	+	0	53
23. —	9	20	49	Paris	-	-	-	-	+	0	40
	10	22	42	Upfalixæ	-	-	-	-	-	0	3*
	10	25	10	Cap. B. Spei	-	-	-	-	+	0	44
25. —	10	46	24	Selenginsk	-	-	-	-	+	1	6*
28. —	16	48	41	Paris	-	-	-	-	-	0	12*
30. —	11	7	58	Grenovici	-	-	-	-	+	0	27
	12	17	43	Upfalixæ	-	-	-	-	+	1	2*
	12	21	32	Cap. B. Spei	-	-	-	-	+	0	28
	12	58	50	Cajaneburgi	-	-	-	-	+	0	15
Sept. 1. —	9	49	40	Inf. Rodrig.	-	-	-	-	+	0	40
8. —	7	55	37	Maffiliae	-	-	-	-	-	0	17*
	9	23	40	Cajaneburgi	-	-	-	-	+	1	1*
15. —	9	39	55	Paris	-	-	-	-	+	0	5
24. Em.	8	7	46	Grenovici	-	-	-	-	+	1	1
	8	29	11	Maffiliae	-	-	-	-	+	0	55
	9	21	35	Cap. B. Spei	-	-	-	-	+	0	47
Oct. 1. —	10	13	56	Paris	-	-	-	-	+	0	59
10. —	6	40	8	Paris	-	-	-	-	+	0	45
17. —	8	36	38	Paris	-	-	-	-	+	0	46
Nov. 9. —	8	51	19	Paris	-	-	-	-	+	1	48*
	10	46	26	Paris	-	-	-	-	+	1	6
18. —	5	27	40	Maffiliae	-	-	-	-	+	0	32*
Dec. 2. —	10	5	4	Stockholmiaæ	-	-	-	-	+	0	32*
	4	33	8	Stockholmiaæ	-	-	-	-	+	0	39
11. —	6	25	44	Stockholmiaæ	-	-	-	-	+	0	43

Observationes secundi satellitis, eodem anno habitæ.

		<sup>h</sup>							
June 11. Im.	14	51	24	Cap. B. Spei	-	+	0	14	
18. —	17	27	11	Cap. B. Spei	-	+	0	4	
July 13. —	13	28	53	Paris	-	-	-	+	0 9
20. —	16	5	1	Paris	-	-	-	+	0 35
	15	54	28	Grenovici	-	-	-	+	1 58*
	17	9	42	Caput B. Spei	-	-	-	+	0 19
31. —	14	57	17	Selenginsk	-	-	-	+	1 41*
Aug. 7. —	10	30	9	Grenovici	-	-	-	+	0 59
	11	43	1	Stockh. optima	-	-	-	+	0 7
	11	43	26	Cap. B. Spei	-	-	-	+	1 17*
14. —	14	21	37	Stockh. certa	-	-	-	+	0 17
	13	18	39	Paris	-	-	-	+	0 25
25. —	12	15	4	Selenginsk	-	-	-	+	0 9
Sept. 1. —	9	1	6	Stockh. certa	-	-	-	—	0 5
	9	2	20	Cap. B. Spei	-	-	-	+	0 16
	8	59	19	Upſaliæ	-	-	-	+	1 2*
8. —	10	37	59	Parifiis	-	-	-	+	0 26
	10	29	30	Grenovici	-	-	-	—	0 5
	10	50	40	Maſſiliæ	-	-	-	—	0 6
	11	42	20	Cap. B. Spei.	-	-	-	+	0 30
15. —	13	6	36	Grenovici	-	-	-	+	3 12*
Oct. 28. Em.	8	9	51	Maſſiliæ	-	-	-	—	0 7
Nov. 11. —	13	12	43	Paris	-	-	-	—	0 2
22. —	5	18	12	Maſſiliæ	-	-	-	+	0 40
	6	9	24	Stockh. dub.	-	-	-	+	0 18*
Dec. 6. —	10	15	59	Paris	-	-	-	+	0 46

Eclipses tertii satellitis Jovis, observatæ anno 1761.

		<sup>h</sup>	<sup>'</sup>	<sup>"</sup>		<sup>'</sup>	<sup>"</sup>
Maj. 11. Im.	15	19	3	Paris	- - - - -	3	30
Jun. 16. Em.	15	26	32	Cap. B. Spei	- - - - -	2	21*
23. Im.	15	20	6	Paris	- - - - -	5	20
	15	32	20	Maffiliæ	- - - - -	5	25
	15	10	24	Grenovici	- - - - -	4	48
Jul. 29. —	11	18	49	Paris	- - - - -	4	10*
	12	23	23	Stockh. certa	- - - - -	5	54
	12	24	0	Cap. B. Spei	- - - - -	5	8
Em.	15	17	13	Cap. B. Spei	+ - - - -	0	23
	15	15	22	Stockh. certa	+ - - - -	0	51
Aug. 5. Im.	15	34	51	Paris	- - - - -	6	18
Sept. 3. —	8	36	36	Cap. B. Spei	- - - - -	5	25
	9	13	39	Cajaneburg	- - - - -	5	4
10. —	11	27	6	Grenovici	- - - - -	4	53
	11	49	9	Maffiliæ	- - - - -	5	37
Nov. 21. Em.	6	41	9	Paris	- - - - -	0	15
	6	52	44	Maffiliæ	- - - - -	0	51
28. Im.	8	10	29	Paris	- - - - -	5	24
	8	23	31	Maffiliæ	- - - - -	6	18
	9	14	22	Stockh. optima	- - - - -	6	28
Em.	10	40	57	Paris	- - - - -	0	16
	10	52	55	Maffiliæ	- - - - -	0	6
	11	44	34	Stockh. dub.	- - - - -	1	4*

Eclipses quarti satellitis, observatæ Anno 1761:

Aug. 10. Im.	<sup>h</sup> 14	<sup>'</sup> 25	<sup>"</sup> 15	Parisiis - - - -	+	<sup>'</sup> 2	<sup>"</sup> 18
	15	32	57	Cap. bonæ Spei —	o	59	
27. —	8	41	40	Grenovici - -	—	o	21
	9	2	3	Maffiliæ - - -	+	o	35
	9	53	32	Cap. B. Spei -	+	1	22
	9	50	18	Upsaliæ - - -	+	1	21
	10	16	23	Tornæ - - - -	+	1	34
Em.	11	47	41	Paris - - - -	+	3	10
	12	1	52	Maffiliæ - - -	+	1	8*
	12	49	o	Cap. B. Spei. -	+	6	16
	12	48	7	Upsaliæ - - -	+	3	54
	12	47	31	Stockh. certa -	+	6	10
	13	15	13	Tornæ - - - -	+	3	6
Nov. 2. Im:	10	51	47	Maffiliæ - - -	+	5	50
Em.	12	57	37	Maffiliæ - - -	+	7	44
19. Im.	6	6	34	Stockh. tub. max. —	8	51	
	6	2	30	Ibid. tub. minore —	4	47	
	4	57	16	Paris - - - - -	—	2	23
Em.	6	56	29	Paris - - - - -	+	6	32

Observationes asterisco \* notatæ, minus certas suspicor. Differentiæ inter reliquas tribuendæ videntur differentiis telescopiorum, majori minorive aëris serenitati, et fortassis cuidam incertitudini circa differentias meridianorum.

Observationes Stockholmenfes habitæ sunt telescopio novo Dollondiano 10 ped. objecta 120<sup>es</sup> secundum diametros amplificante.